

Title	Developing bacteriocins of lactic acid bacteria into next generation biopreservatives
Authors	Field, Des; Ross, R. Paul; Hill, Colin
Publication date	2018-02-15
Original Citation	Field, D., Ross, R. P. and Hill, C. 'Developing bacteriocins of lactic acid bacteria into next generation biopreservatives', Current Opinion in Food Science, In Press. doi: 10.1016/j.cofs.2018.02.004
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://www.sciencedirect.com/science/article/pii/S221479931730070X - 10.1016/j.cofs.2018.02.004
Rights	© 2018 Elsevier Ltd. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. - http://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2023-05-05 09:38:19
Item downloaded from	http://hdl.handle.net/10468/5545

Accepted Manuscript

Title: Developing bacteriocins of lactic acid bacteria into next generation biopreservatives

Author: Des Field R Paul Ross Colin Hill

PII: S2214-7993(17)30070-X

DOI: <https://doi.org/doi:10.1016/j.cofs.2018.02.004>

Reference: COFS 331

To appear in:

Received date: 28-11-2017

Revised date: 6-2-2018

Accepted date: 12-2-2018

Please cite this article as: Field, D., Ross, R.P., Hill, C., Developing bacteriocins of lactic acid bacteria into next generation biopreservatives, *COFS* (2018), <https://doi.org/10.1016/j.cofs.2018.02.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Developing bacteriocins of lactic acid bacteria into next generation biopreservatives

Des Field^{1*}, R Paul Ross¹ and Colin Hill^{1}**

¹ APC Microbiome Institute and School of Microbiology, University College Cork, Western Road, Cork T12 YN60, Ireland.

Key Words: Bacteriocin, lactic acid bacteria, nisin, preservative, antimicrobial peptide, food safety, bioengineering.

*Corresponding author:

*des.field@ucc.ie; Tel: +353-(0)21-4903146; Fax. +353-(0)21-4903101

** c.hill@ucc.ie; Tel. +353-(0)21-4901373; Fax. +353-(0)21-4903101

Running Title: Developing LAB bacteriocins

32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57

Abstract

Bacteriocins are ribosomally synthesized peptides produced by bacteria which can kill other bacteria. Those produced by lactic acid bacteria (LAB) are of great interest as they are often employed in food processing and food fermentations as natural biopreservatives. In this review, we discuss the implementation of bioengineering to enhance the antimicrobial activity, antibacterial spectrum and physico-chemical properties of LAB bacteriocins. Additionally, we discuss the potential applications of bacteriocin derivatives for use as promising food preservatives alone or in combination with other naturally derived antimicrobials as a form of hurdle technology and the regulatory status of strains engineered through food-grade approaches.

Highlights

- Bioengineering can generate novel bacteriocin variants for specific purposes.
- Genome mining has identified new bacteriocin biosynthetic gene clusters.
- Bioengineered bacteriocins show great promise as synergists in hurdle technology.
- Bacteriocin producing strains which have been tailored through food-grade methods can be directly added to food.

Introduction

The growth in world population and the globalization of food commerce has led to large scale food production practices requiring ever longer transport networks and extended storage times until final distribution to consumers. In addition, growing consumer demand for food products that are minimally processed and free from chemical additives presents a complicated and difficult challenge for food processors. Such demand has opened up new opportunities for the use of natural antimicrobials derived from plant, animal or microbial sources to control the growth of undesirable micro-organisms in food [1,2]. Bacteriocins (ribosomally-produced, small, heat-stable peptides that are active against other bacteria) provide one potential solution. While bacteriocins can be produced by a range of Gram-positive and Gram-negative bacteria [3], those produced by lactic acid bacteria (LAB) are of particular interest to the food industry for several reasons. Firstly, members of the LAB group have a history of safe use as starter cultures in food fermentations and many possess “Generally Regarded as Safe (GRAS)” status according to the US Food and Drug

Administration [4]. Secondly, besides being non-toxic to eukaryotic cells, LAB bacteriocins are extremely potent against many food spoilage microbes and pathogenic bacteria, demonstrating killing activity in the nanomolar range. Thirdly, they do not interfere with the sensory quality of foods. Finally, the ribosomal origin of bacteriocins has enabled the manipulation of the associated structural gene in a more direct fashion than is possible for other classes of antimicrobials to obtain variants with potentially beneficial properties. Indeed, several groups have reported on the enhancement of bacteriocins' performance in food environments, including the engineering of derivatives with enhanced activity and inhibition spectra, increased resistance to proteolytic enzymes as well as the combination of such derivatives with other natural antimicrobials in the form of hurdle technology. This review will focus on recent developments with regard to these achievements and present the latest innovations which aim to harness the full potential of these highly potent antimicrobials.

Classification of LAB Bacteriocins

Bacteriocins produced by LAB represent a heterogeneous group of peptides encoded by a diverse genetic repertoire. Some are post-translationally modified and this aspect, together with their mode of action, has traditionally been used as a basis for their classification [5]. The simplest scheme comprises two classes; i.e. Class I bacteriocins also known as RiPPs (Ribosomally Produced and Post-translationally modified Peptides) encompasses all the peptides that undergo enzymatic modification during their biosynthesis (including lanthionines, glycosylation and/or heterocycles). Class II do not contain unusual modifications. However, recent extensive genome mining analysis of LAB suggests that the repertoire of antimicrobials that are encoded in publicly available sequence databases could be even more extensive than previously thought, with some putative classes thus far not reported in LAB (e.g. lasso peptides and sactipeptides) [6]. Furthermore, an *in silico*

screening approach of genome-sequenced isolates from the human gastrointestinal tract (GIT) identified more than 70 clusters of note from almost 60 unique members including Firmicutes, Bacteroidetes, Actinobacteria, and others [7]. The most commonly identified class of bacteriocin was the >10 kDa class, formerly known as bacteriolysins, followed by lantibiotics and sactipeptides [7]. Consequently, a revised scheme (that is also valid for bacteriocins from non-LAB micro-organisms) proposes three classes in which Class I is divided into six subclasses representing different modifications, Class II comprises four subclasses of unmodified peptides of <10kDa and Class III are large-molecular-weight (>10kDa) proteins and are subdivided into the bacteriolysins and the non-lytic bacteriocins [6].

Bioengineering to modulate the physicochemical properties of LAB bacteriocins

Despite the amount of research that has been carried out on the discovery, characterization and mode of action of LAB bacteriocins over the last few decades, to date, just a handful have been commercialized to any extent. These include nisin, a lantibiotic peptide produced by *Lactococcus lactis* [8], pediocin PA-1 produced by *Pediococcus acidilactici* [9] and carnocyclin A produced by *Carnobacterium maltaromaticum* UAL307 [10]. However, studies with other LAB bacteriocins including enterocin AS-48 [11] or lacticin 3147 [12] demonstrate their enormous potential as biopreservatives in food. Nisin is used in most major food-producer countries as a concentrated fermentate powder (e.g. Nisaplin) in a wide variety of dairy and non-dairy products to control the growth of Gram positive bacteria [13]. A fermentate powder produced from the pediocin-producing strain *Pediococcus acidilactici* (ALTA 2351, Kerry Biosciences, Ireland) can be used to protect meat products from *L. monocytogenes* contamination [14]. Carnocyclin A is marketed as Micocin in the US and Canada and has been developed to inhibit *Listeria monocytogenes* in ready-to-eat meat (RTE) products [10]. Accordingly, these bacteriocins have been the subject of several

bioengineering strategies (For comprehensive reviews see [15-17]) that have sought to improve bacteriocin efficacy in the food environment. Peptide function can be influenced by a number of factors including fat content, proteolytic degradation, polar or non-polar food components, pH (which influences the solubility of the bacteriocin) and sodium chloride concentrations [18]. For example, the limited activity spectrum of nisin with respect to pH and its intrinsic insolubility has emphasized the need for alternative versions that exhibit superior stability and are suitable for food fermentation and preservation practices. The natural variant nisin Z, which differs from nisin A by one amino acid (asparagine rather than histidine at position 27) provides an example of a derivative with improved functional characteristics since, while it has similar antimicrobial activity to nisin A, nisin Z displays a higher rate of diffusion [19] and is less soluble at low pH [20]. Recently, bioengineered nisin derivatives were identified with an enhanced ability to diffuse through complex polymers that enabled the peptides to surpass nisin A in restricting growth of *Listeria monocytogenes* in commercially produced chocolate milk containing carrageenan as a stabilizer [21]. Despite the fact that derivatives of the unmodified class II bacteriocins can be generated with relative ease [22], there are comparatively few examples of instances in which LAB producers of the class II bacteriocins have been engineered to positive effect. However, in the case of pediocin PA-1/AcH, greater resistance to oxidation was achieved by the replacement of a methionine moiety with a hydrophobic one, which had only minor effects on antimicrobial activity [23]. Such modifications are an important step in the development of pediocin PA-1 into an advantageous food additive. Some studies have aimed to incorporate protease resistance into LAB-derived peptides. For example, particular modification of trypsin recognition sites in the class IIb bacteriocin salivaricin P produced by *Lactobacillus salivarius* had only minor effects on activity [24].

Bioengineering to modulate the antimicrobial activity and spectrum of LAB bacteriocins

The range of inhibitory activity by LAB bacteriocins can be either narrow, inhibiting only those strains that are closely related to the producer organism, or wide, inhibiting a broad range of Gram-positive micro-organisms [25]. Several investigations have sought to identify nisin derivatives that are enhanced with respect to the purpose for which nisin is most renowned, the inhibition of Gram-positive bacteria [15]. One remarkable derivative, nisin A M21V (Nisin V) (Fig. 1) exhibits enhanced potency against a wide range of targets, most notably *L. monocytogenes*. Furthermore, this enhanced activity was apparent in food model experiments with purified peptide [26]. *L. monocytogenes* is of major concern to the food industry. Apart from the risk to human health, food product recalls due to *Listeria* contamination present an enormous financial burden, estimated to be in the billions of dollars per year in the United States [27]. Recently, nisin V in the form of a fermentate, combined more effectively than nisin A with the essential oils carvacrol, thymol and trans-cinnamaldehyde to inhibit *L. monocytogenes* in a validated food model system [28]. Worryingly, *L. monocytogenes* has the ability to form biofilms providing it with the means to survive on contact surfaces, with higher levels of resistance to disinfectants and potential for growth under the rigorous conditions used for food processing. This could lead to contamination of food products. A recent study demonstrated the effectiveness of nisin M21A, in combination with natural food-grade additives, in targeting biofilms of *L. monocytogenes* F6854 [29], a strain that has been associated with contaminated turkey frankfurters. Similarly, the advantage of shortening or extending the hinge region of nisin has generated variants with improved bioactivity against one or more indicator targets, including *L. monocytogenes*, *E. faecalis* and *B. sporothermodurans* [30]. Bioengineering to enhance the efficacy of Class IIa bacteriocins has also been considered. Mutated peptides in which

residues were substituted within the N-terminal half of pediocin PA-1 exhibited increased activity against *Micrococcus luteus* and *Staphylococcus aureus* [31], while variants within the C-terminus displayed increased potency towards *L. monocytogenes* [32]. Directed mutagenesis studies have been employed to manipulate enterocin AS-48, a broad-spectrum circular bacteriocin produced by a strain of *E. faecalis* [33]. The authors determined that enterocin AS-48 was active in a dimeric form, and established the essential residues involved in these interactions [34]. Such knowledge may facilitate the design and expression of novel variants with improved antimicrobial activity. Notably, the successful chemical synthesis of circular bacteriocins including enterocin AS-48 has recently been achieved [35], providing a more suitable means to generate novel variants, the production of which can often be compromised in the native producer.

The poor activity of LAB bacteriocins toward Gram negative bacteria is due to the outer membrane (OM) of the Gram negative cell wall [36]. The OM functions as an efficient permeability barrier and is able to exclude macromolecules such as bacteriocins or enzymes. Importantly, bioengineered nisin variants S29A and S29G have been shown to display improved activity against Gram negative bacteria [37]. Moreover, while nisin A has been shown to be effective against Gram negatives when used in combination with chelating agents such as EDTA [38], perhaps a more attractive option for food applications is the combination of nisin with natural phytochemical compounds such as essential oils, which act by permeabilization/disruption of the OM. [39]. Recent investigations have also sought to extend the antimicrobial spectrum of pediocin-like bacteriocins to include Gram-negative bacteria. The enterocin CRL35, a pediocin-like bacteriocin, has a potent antilisterial activity but is inactive against Gram-negative targets. In contrast, microcin V (previously known as colicin V) is specifically active against Gram-negative bacteria [40]. A hybrid bacteriocin named Ent35–MccV resulting from the gene fusion of the enterocin CRL35 and microcin V

genes (*munA* and *cvaC*, respectively) displayed inhibitory activity against *E. coli*, *L. monocytogenes*, and other pathogenic Gram-positive and Gram-negative bacteria [41]. Synthetic biology approaches are another promising means to provide insights into structure-stability relationships and generate novel derivatives with improved function. For example, analogues of Lacticin 481 containing non-proteinogenic amino acids were found to have enhanced antibacterial activity [42].

Heterologous expression to increase yield and multiple bacteriocin production

One of the issues that remains to be tackled is that of bacteriocin production, which can be low or inconsistent in lactic acid bacteria and which can be affected by poor growth of producing strains in particular food environments. Although the heterologous production of bacteriocins by LAB is reliant on several factors, bioengineering techniques can facilitate increased levels of bacteriocin production. Some efforts to increase bacteriocin yield have involved using synthetic genes encoding bacteriocins cloned and expressed in yeasts. For instance, the use of codon optimization was recently employed to overcome the bottleneck of low yield of Enterocin A, a class IIa bacteriocin produced by *Enterococcus faecium* CTC492 [43]. Likewise, a recent study involving synthetic biology approaches describes the development of a genetic system that facilitates significant overproduction of nisin [44]. Such innovative systems could potentially reduce the cost of bacteriocin production and also

provides a means by which sufficient quantities of bacteriocin can be produced *in situ*. In addition, novel bacteriocin clusters identified through genome mining and considered highly advantageous to the food industry could be cloned and expressed in suitable hosts. In fact, bacteriocin producing bioprotective cultures that target particular pathogens are commercially available under various trade names [45]. Remarkably, naturally occurring multi-bacteriocin producing LAB have been reported [46]. Consequently, the use of bacteriocin combinations or bacteriocin ‘loading’ may represent a useful approach whereby an assortment of bacteriocins produced *in situ* provide an effective cocktail that can act synergistically to inhibit desired target pathogens. Indeed, some optimistic reports are emerging on the use of multi-bacteriocin mixtures for the effective control of foodborne pathogens such as *Listeria* [47,48].

Bioengineering and the Regulation of Genetically Modified Micro-organisms

Although the application of bioengineering has been instrumental in the fundamental analyses of bacteriocin biology, mode of action studies, and the ability to design more potent peptides with enhanced properties and target selectivity, the application of such peptides as food preservatives may face a significant regulatory obstacle in some jurisdictions. The genetic manipulation of bacteriocins or the producer strains needs to pass through strict safety regulations and guidelines laid down by regulatory agencies such as the FDA (or the European Food Safety Authority [EFSA] in Europe) for approval to be used in human consumption. Indeed, some of the strategies employed to bioengineer many of the bacteriocins described above involve methods that could result in the producer being labelled as a genetically modified micro-organism (GMM). Alternatively, self-cloning of non-pathogenic micro-organisms is not considered to lead to a GMM so long as containment of

the organism is guaranteed (directive 90/219/EC). Notably, the temporary introduction of plasmid vectors or the use of recombinant vectors with an extended history of safe use in the particular micro-organisms, or introduction of DNA from another micro-organism belonging to the same species fall within the definition of self-cloning. Thus, minimal changes to bacteriocin structural genes (such as the alteration of single codons) made using food grade strategies [26,49] fall outside the remit of the EFSA Contained Use legislation and therefore are not regulated as GMMs.

Conclusions

A broad range of technologies have emerged in recent years that provide a battery of valuable tools to expand the potential of bacteriocinogenic strains for food applications. The knowledge gained will improve our understanding on the global effects of bacteriocins in food ecosystems and permit more rational approaches for their application in foods. Several bioengineered bacteriocins capable of inhibiting food-associated Gram positive and Gram-negative bacteria of concern have been recently described. A number have already been tested with satisfactory results in terms of the control of pathogenic bacteria in model food systems. Moreover, their use in combination with other naturally derived antimicrobials in the form of hurdle technology may open new possibilities for the control of a broad range of undesirable organisms. Although genetic manipulation by recombinant and bioengineering-based approaches hold great promise, only bacteriocins which have been tailored through food-grade methodologies can be directly added to food. Furthermore, as the number of microbial genome sequences has increased exponentially, an even larger collection of

putative bacteriocin biosynthetic gene clusters has been revealed. These clusters can be used to identify producer strains, or the information gained from their analysis can be used indirectly to guide the bioengineering of new and existing peptide structures with enhanced functionality for use in the food industry.

Acknowledgements

DF, CH and RR are supported by the Irish Government under the National Development Plan, through SFI Investigator awards to CH and RR (10/IN.1/B3027), and the APC Microbiome Institute under Grant Number SFI/12/RC/2273.

References

1. Lucera A, Costa C, Conte A, Del Nobile MA: **Food applications of natural antimicrobial compounds.** *Front Microbiol* 2012, **3**:287.
2. Gálvez A, Abriouel H, Benomar N, Lucas R: **Microbial antagonists to food-borne pathogens and biocontrol.** *Curr Opin Biotechnol* 2010, **21**:142-148.
3. Riley MA, Wertz JE: **Bacteriocins: Evolution, Ecology, and Application.** *Annual Review of Microbiology* 2002, **56**:117-137.
4. Gálvez A, Abriouel H, López RL, Omar NB: **Bacteriocin-based strategies for food biopreservation.** *International Journal of Food Microbiology* 2007, **120**:51-70.
5. Cotter PD, Ross RP, Hill C: **Bacteriocins - a viable alternative to antibiotics?** *Nat Rev Microbiol* 2013, **11**:95-105.
6. Alvarez-Sieiro P, Montalban-Lopez M, Mu D, Kuipers OP: **Bacteriocins of lactic acid bacteria: extending the family.** *Appl Microbiol Biotechnol* 2016, **100**:2939-2951.

7. Walsh CJ, Guinane CM, Hill C, Ross RP, O'Toole PW, Cotter PD: **In silico identification of bacteriocin gene clusters in the gastrointestinal tract, based on the Human Microbiome Project's reference genome database.** *BMC Microbiology* 2015, **15**:183.
8. Delves-Broughton J, Blackburn P, Evans RJ, Hugenholtz J: **Applications of the bacteriocin, nisin.** *Antonie Van Leeuwenhoek* 1996, **69**:193-202.
9. Papagianni M, Anastasiadou S: **Pediocins: The bacteriocins of Pediococci. Sources, production, properties and applications.** *Microb Cell Fact* 2009, **8**:3.
10. Martin-Visscher LA, Yoganathan S, Sit CS, Lohans CT, Vederas JC: **The activity of bacteriocins from Carnobacterium maltaromaticum UAL307 against Gram-negative bacteria in combination with EDTA treatment.** *FEMS Microbiology Letters* 2011, **317**:152-159.
11. Sanchez-Hidalgo M, Montalban-Lopez M, Cebrian R, Valdivia E, Martinez-Bueno M, Maqueda M: **AS-48 bacteriocin: close to perfection.** *Cell Mol Life Sci* 2011, **68**:2845-2857.
12. Suda S, Cotter PD, Hill C, Ross RP: **Lacticin 3147--biosynthesis, molecular analysis, immunity, bioengineering and applications.** *Curr Protein Pept Sci* 2012, **13**:193-204.
13. Gharsallaoui A, Oulahal N, Joly C, Degraeve P: **Nisin as a Food Preservative: Part 1: Physicochemical Properties, Antimicrobial Activity, and Main Uses.** *Crit Rev Food Sci Nutr* 2016, **56**:1262-1274.
14. López-Cuellar MdR, Rodríguez-Hernández A-I, Chavarría-Hernández N: **LAB bacteriocin applications in the last decade.** *Biotechnology & Biotechnological Equipment* 2016, **30**:1039-1050.
- A comprehensive review on recent LAB bacteriocin applications and their role in safety, quality and improvement of human health.
15. Field D, Cotter PD, Ross RP, Hill C: **Bioengineering of the model lantibiotic nisin.** *Bioengineered* 2015, **6**:187-192.
16. Ramu R, Shirahatti PS, Devi AT, Prasad A, Kumuda J, Lachana MS, Zameer F, Dhananjaya BL, Nagendra Prasad MN: **Bacteriocins and Their Applications in Food Preservation.** *Crit Rev Food Sci Nutr* 2015: DOI: 10.1080/10408398.2015.1020918
17. Porto MC, Kuniyoshi TM, Azevedo PO, Vitolo M, Oliveira RP: **Pediococcus spp.: An important genus of lactic acid bacteria and pediocin producers.** *Biotechnol Adv* 2017, **35**:361-374.
18. Mills S, Ross RP, Hill C: **Bacteriocins and bacteriophage; a narrow-minded approach to food and gut microbiology.** *FEMS Microbiol Rev* 2017, **41**:S129-s153.
- **This is a comprehensive review on the use and potential of bacteriocins and peptides in food systems.
19. de Vos WM, Mulders JW, Siezen RJ, Hugenholtz J, Kuipers OP: **Properties of nisin Z and distribution of its gene, nisZ, in Lactococcus lactis.** *Appl Environ Microbiol* 1993, **59**:213-218.
20. Rollema HS, Kuipers OP, Both P, de Vos WM, Siezen RJ: **Improvement of solubility and stability of the antimicrobial peptide nisin by protein engineering.** *Appl Environ Microbiol* 1995, **61**:2873-2878.
21. Rouse S, Field D, Daly KM, O'Connor PM, Cotter PD, Hill C, Ross RP: **Bioengineered nisin derivatives with enhanced activity in complex matrices.** *Microb Biotechnol* 2012, **5**:501-508.
22. O'Shea EF, Cotter PD, Ross RP, Hill C: **Strategies to improve the bacteriocin protection provided by lactic acid bacteria.** *Curr Opin Biotechnol* 2013, **24**:130-134.
23. Kaur K, Andrew LC, Wishart DS, Vederas JC: **Dynamic relationships among type IIa bacteriocins: temperature effects on antimicrobial activity and on structure of the C-terminal amphipathic alpha helix as a receptor-binding region.** *Biochemistry* 2004, **43**:9009-9020.
24. O'Shea EF, O'Connor PM, Cotter PD, Ross RP, Hill C: **Synthesis of trypsin-resistant variants of the Listeria-active bacteriocin salivaricin P.** *Appl Environ Microbiol* 2010, **76**:5356-5362.
25. Klaenhammer TR: **Bacteriocins of lactic acid bacteria.** *Biochimie* 1988, **70**:337-349.
26. Field D, Quigley L, O'Connor PM, Rea MC, Daly K, Cotter PD, Hill C, Ross RP: **Studies with bioengineered Nisin peptides highlight the broad-spectrum potency of Nisin V.** *Microb Biotechnol* 2010, **3**:473-486.

27. Thomsen MR, McKenzie AM: **Market Incentives for Safe Foods: An Examination of Shareholder Losses from Meat and Poultry Recalls.** *American Journal of Agricultural Economics* 2001, **83**:526-538.
28. Field D, Daly K, O'Connor PM, Cotter PD, Hill C, Ross RP: **Efficacies of nisin A and nisin V semipurified preparations alone and in combination with plant essential oils for controlling *Listeria monocytogenes*.** *Appl Environ Microbiol* 2015, **81**:2762-2769.
29. Smith MK, Draper LA, Hazelhoff PJ, Cotter PD, Ross RP, Hill C: **A Bioengineered Nisin Derivative, M21A, in Combination with Food Grade Additives Eradicates Biofilms of *Listeria monocytogenes*.** *Front Microbiol* 2016, **7**:1939.
30. Zhou L, van Heel AJ, Montalban-Lopez M, Kuipers OP: **Potentiating the Activity of Nisin against *Escherichia coli*.** *Frontiers in Cell and Developmental Biology* 2016, **4**:7.
- This study demonstrates how combining nisin with fragments of eukaryotic antimicrobial peptides can improve the activity of lantibiotics against Gram-negative bacteria.
31. Song DF, Li X, Zhang YH, Zhu MY, Gu Q: **Mutational analysis of positively charged residues in the N-terminal region of the class IIa bacteriocin pediocin PA-1.** *Lett Appl Microbiol* 2014, **58**:356-361.
32. Sun L, Song H, Zheng W: **Improvement of Antimicrobial Activity of Pediocin PA-1 by Site-directed Mutagenesis in C-terminal Domain.** *Protein Pept Lett* 2015, **22**:1007-1012.
33. Maqueda M, Galvez A, Bueno MM, Sanchez-Barrena MJ, Gonzalez C, Albert A, Rico M, Valdivia E: **Peptide AS-48: prototype of a new class of cyclic bacteriocins.** *Curr Protein Pept Sci* 2004, **5**:399-416.
34. Cebrian R, Martinez-Bueno M, Valdivia E, Albert A, Maqueda M, Sanchez-Barrena MJ: **The bacteriocin AS-48 requires dimer dissociation followed by hydrophobic interactions with the membrane for antibacterial activity.** *J Struct Biol* 2015, **190**:162-172.
35. Hemu X, Qiu Y, Nguyen GK, Tam JP: **Total Synthesis of Circular Bacteriocins by Butelase 1.** *J Am Chem Soc* 2016, **138**:6968-6971.
36. Nikaido H, Vaara M: **Molecular basis of bacterial outer membrane permeability.** *Microbiol Rev* 1985, **49**:1-32.
37. Field D, Begley M, O'Connor PM, Daly KM, Hugenholtz F, Cotter PD, Hill C, Ross RP: **Bioengineered Nisin A Derivatives with Enhanced Activity against Both Gram Positive and Gram Negative Pathogens.** *PLoS ONE* 2012, **7**:e46884.
38. Brannen JK, Davidson PM: **Enhancement of nisin, lysozyme, and monolaurin antimicrobial activities by ethylenediaminetetraacetic acid and lactoferrin.** *Int J Food Microbiol* 2004, **90**:63-74.
39. Campion A, Morrissey R, Field D, Cotter PD, Hill C, Ross RP: **Use of enhanced nisin derivatives in combination with food-grade oils or citric acid to control *Cronobacter sakazakii* and *Escherichia coli* O157:H7.** *Food Microbiol* 2017, **65**:254-263.
40. Gillor O, Kirkup BC, Riley MA: **Colicins and Microcins: The Next Generation Antimicrobials.** In *Advances in Applied Microbiology*. Academic Press; 2004:129-146.
41. Acuña L, Picariello G, Sesma F, Morero RD, Bellomio A: **A new hybrid bacteriocin, Ent35-MccV, displays antimicrobial activity against pathogenic Gram-positive and Gram-negative bacteria.** *FEBS Open Bio* 2012, **2**:12-19.
42. Knerr PJ, Oman TJ, Garcia De Gonzalo CV, Lupoli TJ, Walker S, van der Donk WA: **Non-proteinogenic amino acids in lactacin 481 analogues result in more potent inhibition of peptidoglycan transglycosylation.** *ACS Chem Biol* 2012, **7**:1791-1795.
43. Hu X, Mao R, Zhang Y, Teng D, Wang X, Xi D, Huang J, Wang J: **Biotechnical paving of recombinant enterocin A as the candidate of anti-*Listeria* agent.** *BMC Microbiol* 2014, **14**:220.
44. Kong W, Lu T: **Cloning and optimization of a nisin biosynthesis pathway for bacteriocin harvest.** *ACS Synth Biol* 2014, **3**:439-445.

- 412 45. Chikindas ML, Weeks R, Drider D, Chistyakov VA, Dicks LM: **Functions and emerging applications**
413 **of bacteriocins.** *Curr Opin Biotechnol* 2017, **49**:23-28.
- 414 *Excellent review on potential applications of bacteriocins and their involvement in the regulation of
415 various processes within microbial communities.
- 416 46. Himeno K, Fujita K, Zendo T, Wilaipun P, Ishibashi N, Masuda Y, Yoneyama F, Leelawatcharamas
417 V, Nakayama J, Sonomoto K: **Identification of enterocin NKR-5-3C, a novel class IIa**
418 **bacteriocin produced by a multiple bacteriocin producer, *Enterococcus faecium* NKR-5-3.**
419 *Biosci Biotechnol Biochem* 2012, **76**:1245-1247.
- 420 47. Vijayakumar P, Muriana P: **Inhibition of *Listeria monocytogenes* on Ready-to-Eat Meats Using**
421 **Bacteriocin Mixtures Based on Mode-of-Action.** *Foods* 2017, **6**:22.
- 422 48. Mills S, Griffin C, O'Connor PM, Serrano LM, Meijer WC, Hill C, Ross RP: **A Multibacteriocin**
423 **Cheese Starter System, Comprising Nisin and Lacticin 3147 in *Lactococcus lactis*, in**
424 **Combination with Plantaricin from *Lactobacillus plantarum*.** *Applied and Environmental*
425 *Microbiology* 2017, **83**.
- 426 *This study demonstrates the potential of using multiple bacteriocin producing strains as starters in
427 laboratory-scale cheeses to effectively control *Listeria* growth.
- 428 49. Cotter PD, Hill C, Ross RP: **A food-grade approach for functional analysis and modification of**
429 **native plasmids in *Lactococcus lactis*.** *Appl Environ Microbiol* 2003, **69**:702-706.




Figure 1. Bioengineering bacteriocins: bacteriocins with improved physicochemical properties (pH, solubility) have been generated as well as novel variants with improved activity, target spectrum and anti-biofilm efficacy. Bioengineered variants can act synergistically to inhibit desired target pathogens in the form of hurdle technology. Genome mining has identified an even larger collection of new bacteriocin biosynthetic gene clusters which can be used to guide the bioengineering of new and existing peptide structures. Bioengineered strains which have been tailored through food-grade approaches are not considered GMMs and can be directly added to food.